

Ethnobotanical usage of plants as aphrodisiac agents in Anatolian folk medicine

Selin Tufan, Gizem Gülsoy Toplan *, Afife Mat

Department of Pharmacognosy, Faculty of Pharmacy, İstanbul University, 34116 İstanbul, Turkey

* Correspondence: eczgizemgulsoy@gmail.com, gizem.toplan@istanbul.edu.tr (G.G.T.); Tel: +90 212 440 00 00; ORCID No: 0000-0002-0544-2532.

Submitted: 24 June 2017 / Revised: 08 August 2017 / Accepted: 14 August 2017

ABSTRACT: Several plants have been used as aphrodisiac to treat sexual problems for centuries. Nowadays, despite the synthesis of chemicals, herbal medicines continue to maintain their importance. Turkey has a great tradition of folk medicine due to its rich flora and many plants have been used for the treatment of several disorders. This work presents aphrodisiac plants traditionally used in Turkey, with their local names, used parts, usage and administration route.

KEYWORDS: Aphrodisiac plants; folk medicine; ethnobotanical usage; aphrodisiacs

1. INTRODUCTION

The word aphrodisiac is originated from Aphrodite, The Greek Goddess of love and beauty [1]. With all respect to her duty she is responsible from the lovely ways of creating human relations. She is spreading love potions all over the humans and she has been decorating nature with colors with the help of trees and flowers. According to mythology, when Aphrodite born from seafoam and set foot on earth face, different types of flowers started to blossom under her foot and since that day, followers of Aphrodite have been able to use those flowers (which are called "Plants of Love") as aphrodisiac [2-4].

Since the beginning of humankind, sexuality has been really important in terms of continuation of the race and transfer of the culture to the offsprings. The usage of aphrodisiacs has been developed during centuries. Some of them provide psychological relief, while others affect physical system to treat sexual disorders [5-7].

Sexual relationship is very important in human life and it influences the social relationship between partners [1]. The cause of sexual disorders can be physical or psychological, and sometimes both. Today, both of them are lying under the majority of problems [8]. Nowadays, chronic disorders, antipsychotic and antidepressant therapeutic drugs, tobacco, stress, chronic alcohol abuse have increased these problems. The most common sexual problems in men are ejaculation disorders, erectile dysfunction, and loss of libido. In 25% to 63% of women, sexual dysfunction is also a serious problem [6-8].

By definition, aphrodisiacs are materials that are stimulating libido [9]. Regarding to different perspectives, their perception can be variable. During history, aphrodisiac agents have been used as stimulant of sexual intercourse, pleasure booster, flaming agents of libido and an agent to strip from shyness. Different countries from all around the world, have formed their own culture with various plants, according to the knowledge transferred from their ancestors. They adapted those plants into their traditional medicinal system [4]. There are different ways to use them, such as cigarette, tea, incense, pill, spice or meal, ointment and oil. In folkloric medicine of Chinese, Indian, Egyptian, Roman, Anatolian and Greek cultures, there are many different herbal and animal remedies [4-10]. The most common used plants or drugs are *Cannabis sativa*, opium, *Panax ginseng*, *Ginkgo biloba*, *Zingiber officinale*, *Epimedium sp.*, *Mucuna pruriens*, *Corynanthe yohimbe*, *Chelidonium majus*, *Tribulus terrestris*, *Crocus sativus*, *Phoenix dactylifera*, *Pegalum harmala*, *Catha edulis*, *Salvia haematodes*, *Lepidium meyenii*, *Asparagus racemosus*, *Glycyrrhiza glabra*. Various foods also have been used as aphrodisiac agent such as vegetables [artichokes, asparagus], spices (anise, basil, coriander, fennel, sage, ginger, garlic, and saffron), oysters, chocolate, and fruits [10-14].

How to cite this article: Tufan S, Gülsoy Toplan G, Mat A. Ethnobotanical usage of plants as aphrodisiac agents in Anatolian folk medicine. Marmara Pharm J. 2018; 22 (2): 142-151.

There are several ethnopharmacological reports on the medicinal uses of aphrodisiac plants of various cultures. Some of these plants or animal products were investigated for their aphrodisiac activity. The potential aphrodisiac properties were examined using *in vitro* assays, different animal models or clinical trials [11-14]. Although most studies demonstrated positive aphrodisiac activities, further studies are needed to reveal their mechanism of activity.

Ambrein and *Panax ginseng* have been shown to relax corpus cavernous smooth muscle or other smooth muscles in animal models, *in vitro* assays [15-16]. Hydro-alcoholic and aqueous extracts of *Asparagus racemosus* (400 mg/kg body weight) have showed aphrodisiac properties on male wistar albino rats and these extracts have increased number of mounts and mating performance, and lower dose (200mg/kg body weight) of hydro-alcoholic extract have shown moderate activity. In addition, 200 mg/kg body weight of milk decoction of *A. racemosus* roots has shown significant aphrodisiac effect. It increased mount frequency and caused decrease in mount latency, post ejaculatory latency and ejaculation latency [17]. There is one study showing that aqueous extract of *Glycyrrhiza glabra* roots and rhizomes have significant decreasing effects on mount latency and intromission latency when 150 mg/kg & 300 mg/kg body weight/day administrated orally by gavage for 28 days on sexually active male rats [18]. *Tribulus terrestris*, also known as "ghokhru" and its extracts have been used as medicine traditionally in both Asia and India and also in Turkey to treat urinary, cardiovascular and gastrointestinal disorders. It is also, used by humankind to stimulate sexual desires due to its steroidal saponin protodioscin [11-19]. *T. terrestris* has been shown to increase erection quality in animals by observing an increased intracavernous pressure, while yohimbine, ginseng, and saffron increased erection quality in humans [11-20]. Saffron is one of the most expensive and highly valued spices in the world. The effect of *Crocus sativus* (saffron) on male erectile dysfunction was studied for 10 days, researchers followed up 20 male patients with erectile dysfunction whereby they were administered a tablet containing 200 mg of saffron every morning. The selected patients were subjected to the nocturnal penile tumescence (NPT) test and the international index of erectile function questionnaire (IIEF-15) at the beginning and the end of the 10 days treatment. The results showed a statistically significant improvement after 10 days of taking saffron [21-22]. *Passiflora incarnata* has been used as anxiolytic, sedative, anticonvulsant all around the world. According to Dhawan et al. studies, the methanolic extract of *Passiflora incarnata* leaves showed strong aphrodisiac effects [23].

Besides, some of plants are usually used as aphrodisiac agents due to their strong antioxidant activities. Oxidative stress or excessive production of free radicals can damage sexual hormones and also decreased libido. This damage can cause infertility and decrease libido in both women and men. Recently, the studies revealed that lipid peroxidation plays significant role at infertility especially affecting sexual hormones, sperm motility and viability. In order to understand the relation between antioxidant and aphrodisiac activities, the effects of many plant extracts were examined by several researchers. According to these studies, there is a strong correlation between plants antioxidant properties and aphrodisiac activities [24].

Several plants have been used traditionally for centuries in Turkey for the treatment of various sexual disorders. Most of the remedies are prepared from a mixture of several species while there are some strong remedies with only one component.

The most popular traditional aphrodisiac is '*Mesir Macunu*' (Mesir Paste). It is a traditional Turkish gum like candy produced in Manisa (Aegean region of Turkey). The original Mesir paste, prepared from 41 different spices, sugar and honey, is a tradition in Anatolia since 16th century [25]. The majority of historians claims its similarities with theriac. The preparation of *mesir macunu* and its ceremonial distribution are still alive in Manisa [26, 27]. Today the content of mesir paste, given in Table 1, has minor differences from the original one. When the Mesir Paste is examined, it is clearly seen that has high antioxidant activity [18-28-29-30-31-32]. The consist of Mesir paste includes many plants which possess strong antioxidant activity There are several studies on these plants antioxidant activities. Some these studies are given in Table 1.

Table 1. Mesir paste.

Local Name	Plant	Family	Part Used	References	Antioxidant
Anason	<i>Pimpinella anisum</i> L.	Umbelliferae	Ripe Fruits	[11-33-34]	[28-35]
Çivitotu	<i>Isatis tinctoria</i> L.	Fabaceae	Leaves	[33]	[29-36]
Çöpçini / Çin Saparması	<i>Smilax china</i> L.	Liliaceae	Roots	[33]	[37-38]
Çörekotu	<i>Nigella sativa</i> L.	Ranunculaceae	Sun Dried Seeds	[33]	[39-40]
Darifüfül	<i>Piper longum</i> L.	Piperaceae	Bitter Fruits	[33]	[41-42]
Defne	<i>Laurus nobilis</i> L.	Lauraceae	Leaves	[33]	[43-44]
Galanya / Kalanga / Büyük Havlican	<i>Alpinia galangal</i> L.	Zingiberaceae	Rhizomes	[33]	[45-46]
Hindistan Cevizi/Kopra	<i>Cocus nucifera</i> L.	Arecaceae	Dried Seeds	[33-47]	[48]
Hindistan Çiçeği	<i>Myristica fragrans</i> Houutt.	Myristicaceae	Flowers	[12-14-33-49]	[50-51]
Hıyarşenbe	<i>Cassia fistula</i> L.	Fabaceae	Ripe Fruits	[33]	[52-53]
Kakule	<i>Elettaria cardamomum</i> L.	Zingiberaceae	Dried Unripe Fruits	[33]	[54-55]
Kara Halile	<i>Terminalia chebula</i> Retz.	Combretaceae	Fruits	[33]	[56-67]
Karabiber	<i>Piper nigrum</i> L.	Piperaceae	Dried Unripe Fruits	[33]	[58-59]
Karanfil	<i>Syzygium aromaticum</i> L.	Myrtaceae	Flowers	[11-12-33-49]	[60-61]
Kebabiyi / Kebabe / Kübabe	<i>Piper cubeba</i> L.	Piperaceae	Dried Unripe Fruits	[33]	[30]
Kimyon	<i>Cuminum cyminum</i> L.	Umbelliferae	Dried Ripe Fruits	[33]	[62-63]
Kışniş	<i>Coriandrum sativum</i> L.	Umbelliferae	Fruits	[11-33-34]	[64-65]
Küçük Havlican	<i>Galanga officinalis</i> (L.) Wild.	Zingiberaceae	Dried Rhizomes	[33]	[-]
Mercanköşk	<i>Origanum majorana</i> L.	Labiatae	Aerial parts	[33]	[66-67]
Meyan	<i>Glycyrrhiza glabra</i> L.	Leguminosae	Condensed Root Extract	[33]	[31-68]
Mirsafi	<i>Commiphora myrrha</i> (Nees) Engl.	Burceraceae	Usare	[33]	[32-69]
Portakal	<i>Citrus aurantium</i> var. <i>dulce</i> L.	Rutaceae	Pericarpium	[33]	[-]
Ravent	<i>Rheum officinale</i> L.	Polygonaceae	Rhizomes	[33]	[70]
Rezene	<i>Foeniculum vulgare</i> Mill.	Umbelliferae	Dried Ripe Fruits	[11-33]	[71-72]
Safran	<i>Crocus sativus</i> L.	Iridaceae	Stigma	[11-12-14-33]	[73-74]
Sakız	<i>Pistacia lentiscus</i> L.	Anacardiaceae	Resin	[33]	[75-76]
Sarı Halile	<i>Terminalia citrina</i> (Gaertn.) Roxb. ex Fleming	Combretaceae	Fruits	[33]	[77]
Sinameki	<i>Cassia acutifolia</i> Mill.	Leguminosae	Dried Leaves	[33]	[78]
Siyah Hardal	<i>Brassica nigra</i> L.	Cruciferae	Seeds	[33]	[79]
Tarçın	<i>Cinnamomum cassia</i> (L.) J.Presl.	Lauraceae	Cortex	[33]	[80-81]
Tarçın Çiçeği	<i>Cinnamomum zeylanicum</i> L.	Lauraceae	Flowers	[33]	[82]
Teke mersini	<i>Vaccinium myrtillus</i> L.	Ericaceae	Fruits	[33]	[83]
Topalak	<i>Cyclamen coum</i> Miller	Primulaceae	Tubers	[33]	[-]
Udülkahr / Nezeleotu	<i>Anacyclus pyrethrum</i> L.	Asteraceae	Dried Roots	[33]	[-]
Vanilya	<i>Vanilla planifolia</i> Andr.	Orchidaceae	Fermented Fruit	[33]	[84-85]
Yenibahar	<i>Pimenta officinalis</i> L.	Myrtaceae	Dried Unripe Fruits	[33]	[-]
Zencefil	<i>Zingiber officinale</i> Rosc.	Zingiberaceae	Rhizomes	[11-33]	[86-87]
Zerdeçal / Hintsafra / Sarıboya	<i>Curcuma longa</i> L.	Zingiberaceae	Rhizomes	[33]	[88-89]
Zulumba	<i>Curcuma zedoaria</i> Rosc.	Zingiberaceae	Rhizomes	[33]	[90]

2. MATERIALS AND METHODS

Ethnobotanical studies and thesis about Anatolian folk medicine were reviewed and plants used as aphrodisiac were selected. Information about them were reported. Among all studies, the plants used as aphrodisiac agent in more than one publication have been selected and added to the Table 2.

Table 2. The list of Turkish aphrodisiac plants.

Plant	Local Name	Family	Part Used	Preparation	Dosage	References
<i>Prangos ferulacea</i> L.	Çağsır, köfteotu, kürdanotu, melekotu, pıtrak	Apiaceae	Roots	Mixed with sweets		[91]
<i>Prangos meliocarpoides</i> Boiss. var. <i>meliocarpoides</i>	Çarsır, hiltıl, sultan teresi	Apiaceae	Roots	Mixed with sweets		[91]
<i>Morina persica</i> L.	Boğa dikeneni	Morinaceae	Aerial parts, Roots	Eaten raw		[91]
<i>Hippophae rhamnoides</i> L. subsp. <i>caucasica</i>	Pişot	Elaeagnaceae	Fruits	Eaten raw		[92]
<i>Brassica oleracea</i> L. var. <i>capitata</i> L.	Başlı lahana	Cruciferae	Seeds			[93]
<i>Corylus avellana</i> L. var. <i>avellana</i> <i>Corylus colurna</i> L. <i>Corylus maxima</i> Miller	Fındık	Corylaceae	Seeds	1 kg of hazelnut, butter, honey, raisins and 0.5 kg of garlic are pounded and mixed.	Mixture eaten every morning on empty stomach.	[93-94]
<i>Junierus drupacea</i> L.	Andız	Cupressaceae	Cortex	Decoction, Powder		[95]
<i>Orchis spp.</i> L.	Salep	Orchidaceae	Tubers	Powder		[95]
<i>Cerasus mahaleb</i> L.	Mahlep	Rosaceae	Seeds	Powder, Infusion		[95-96]
<i>Capparis spinosa</i> L. var. <i>spinosa</i>	Gebere	Capparaceae	Seeds	Crushed Extract		[97]
<i>Matricaria chamomilla</i> L.	Papatya, deli papatya, tellipapatya	Asteraceae	Capitulum	Infusion		[97]
<i>Rosa canina</i> L.	Kuşburnu, itburnu	Rosaceae	Fruits, Roots, Seeds	Decoction, Fruits (eaten by chewing), Raw, Mixture (jam etc.)	Herbal tea with hibiscus 1-2 glass per day for 2-3 weeks. Paste prepared from fresh fruits (2-3 kg) and paste mixed with honey in the ratio of 1:3. Consumed on empty stomach.	[98-99-100-101-102-103]
<i>Ceratonia siliqua</i> L.	Keçiboynuzu, Harnup	Fabaceae	Fruits	Mixture (jam etc.), Fruits (eaten by chewing), Raw, Decoction, Fruit powder, Spice	2-3 pieces or 2-3 tablespoons are eaten before breakfast for 2-3 weeks.	[98]

Table 3 (Continued). The list of Turkish aphrodisiac plants.

Plant	Local Name	Family	Part Used	Preparation	Dosage	References
<i>Ferula elaeochytris</i> Korovin	Çarşır, çaaşır, çarşır otu, çakşır, çarşır göbeği	Apiaceae	Roots, Flowers, Seeds, Fruits	Spice, Rhizome juice, Fruits (eaten by chewing), Powdered roots	Drunken 1 teacup twice a day for 2-3 weeks. Peeled roots were dried, pulverized and mixed with honey.	[96-98-104]
<i>Gundelia tournefortii</i> L. var. <i>tournefortii</i> L.	Kenger, Hüznbol	Asteraceae	Latex, Roots	A piece of latex is eaten, Decoction		[100-103-105]
<i>Zea mays</i> L.	Mısır	Poaceae	Style	Decoction, Infusion		[100-105-106]
<i>Ferula communis</i> L.	Çarşır, çarşır otu, çakşır, çarşır göbeği	Apiaceae	Roots	Mash	1 teaspoon twice a day for 1-2 weeks	[107]
<i>Opuntia ficus-indica</i> L.	Kaynana dili, Frenk inciri	Cactaceae	Fruits	Peeled and eaten raw		[105]
<i>Elaeagnus angustifolia</i> L.	İğde	Elaeagnaceae	Flowers	Scent		[105]
<i>Calicotome villosa</i> (Poir.) Link	Azgan	Fabaceae	Flowers, Essential oil	Scent		[105]
<i>Sideritis syriaca</i> L. subsp. <i>Nusairiensis</i>	Adaçayı	Lamiaceae	Aerial parts	Decoction		[105]
<i>Sesamum indicum</i> L.	Susam	Pedaliaceae	Seeds	Eaten raw		[105]
<i>Pinus pinea</i> L.	Çam fıstığı	Pinaceae	Seeds, Fruits	Eaten raw		[103-108]
<i>Arceuthos drupacea</i> Ant. et Kotschy.	Andız	Cupressacaceae	Grape molasses			[108]
<i>Urtica dioica</i> L.	Isırgan	Urticaceae	Aerial parts, Seeds, Leaves	Infusion, Crushed, Leaves mixed with sweets		[91]
<i>Melilotus officinalis</i> L.	Kokulu yonca	Fabaceae	Aerial parts	Decoction		[92]
<i>Punica granatum</i> L.	Nar	Punicaceae	Seeds	Eaten raw		[109]

3. RESULTS AND DISCUSSION

According to ethnobotanical studies, 27 medicinal plants are listed as aphrodisiacs in Turkish traditional medicine. These plants are given in Table 2. The most popular plant is *Rosa canina* which is widely used through all Anatolia. The majority of the plants are prepared in the form of infusion or decoction from freshly collected plant parts.

Mesir paste is remarkably the famous traditional aphrodisiac since centuries.

While comparing usage methods of aphrodisiacs in Turkey and other countries, similarities and differences were observed. For example, *Rosa canina* fruits mixed with honey are used in Turkey while in other countries its essential oil is reputed as aphrodisiac. The flowers of *Elaeagnus angustifolia* are recognized as aphrodisiac in Turkish ethnobotanical studies in Iran its fruits are used as female aphrodisiac. *Sesamum indicum* seeds are used against impotency in both Turkey and India. Turkish people prefer eating raw seeds for treatment. Ayurvedic system suggests a mixture of seeds with honey once a day, before meal. *Urtica dioica*, a component of Swiss love potion, is also used in sweet mixtures in Turkey for the same purpose. *Brassica oleracea*, *Corylus avellana*, *Capparis spinosa* have similar aphrodisiac usage in different countries. Although *Lepidium meyenii* is quite popular as aphrodisiac plants in other countries, *Lepidium sativum* have been used as aphrodisiac agent only in Capadocia region from Turkey [110].

The adverse effects and expensive price of synthetic drugs orient the humankind to natural treatment. Natural products have been used for centuries and investigations proved the effectiveness of some of them.

Author contributions: Concept – S.T.,G.G.T.; Design – S.T.,G.G.T., A.M.; Supervision – A.M.; Resource – S.T., G.G.T.; Materials – S.T., G.G.T.; Data Collection and/or Processing S.T.,G.G.T.; Analysis and/or Interpretation – S.T.,G.G.T.,A.M.; Literature Search – S.T.,G.G.T.; Writing – S.T.,G.G.T. A.M.; Critical Reviews – S.T.,G.G.T., A.M.

Conflict of interest statement: The authors declared no conflict of interest.

REFERENCES

- [1] Chauhan NS, Sharma V, Dixit VK, Thakur M. A review on plants used for improvement of sexual performance and virility. *Biomed Res Int.* 2014; 2014: 1-19.
- [2] Vernant JP. *Evren, Tanrılar, İnsanlar.* Dost Kitabevi Yayınları, Ankara. 2001.
- [3] Daly KN. *Greek and Roman Mythology A to Z.* Chelsea House Publishers, New York. 2009.
- [4] Rättsch C. *Plants of Love: Aphrodisiacs in Myth, History, and the Present.* Ten Speed Press, Berkeley. 1997.
- [5] Babacan SS. İnsan Seksüalitesinin Kültürel ve Psikososyal Yönleri. *Kefdergi.* 2003; 11: 131-136.
- [6] İncesu C. Cinsel İşlevler ve Cinsel İşlev Bozuklukları. *J Clin Psy.* 2004; 7(3): 3-13.
- [7] Vural AG. Evlilik öncesi cinsel danışmanlık programının yeni evli çiftlerin cinsel doyumuna etkisi. T.C. Ege Üniversitesi Sağlık Bilimleri Enstitüsü, İzmir. 2007.
- [8] Sadock BJ. *Normal İnsan Cinselliği, Cinsel İşlev ve Cinsel Kimlik Bozuklukları.* Güneş, Ankara. 2007.
- [9] Mathur, M. Herbal aphrodisiac their need, biology and status: Global and regional scenario. *J Nat Prod.* 2012; 5: 131-146.
- [10] Rättsch C, Müller-Ebeling C. *The Encyclopedia of Aphrodisiacs Psychoactive Substances for Use in Sexual Practices.* Park Street Press, Toronto. 2003.
- [11] Melnyk JP, Marccone MF. Aphrodisiacs from plant and animal sources – A review of current scientific literature. *Food Res Int.* 2011; 4: 840-850.
- [12] Sumalatha K, Kumar SA, Lakshmi SM. Review on natural aphrodisiac potentials to treat sexual dysfunction. *IJPT.* 2010; 1(1): 6-14.
- [13] Thakur M, Chauhan NS, Bhargava A, Dixit VK. A comparative study on aphrodisiac activity of some Ayurvedic herbs in male albino rats. *Arch Sex Behav.* 2009; 38 (6): 1009-1015.
- [14] Singh R, Ali A, Gupta G, Semwal A, Jeyabalan G. Some medicinal plants with aphrodisiac potential: A current status. *J Acute Dis.* 2013; 2(3): 179-188.
- [15] Taha SA, Raza M, El-Khawad IE. Effect of ambrein on smooth muscle responses to various agonists. *J Ethnopharmacol.* 1998; 60(1): 19–26.
- [16] Kim HJ, Woo DS, Lee G, Kim, JJ. The relaxation effects of ginseng saponin in rabbit corporal smooth muscle: Is it a nitric oxide donor? *Br J Urol.* 1998; 82(5): 744–748.
- [17] Wani JA, Achur RN, Nema RK. Phytochemical screening and aphrodisiac property of *Asparagus racemosus*. *Int J Pharm Drug Res.* 2011; 3(2): 112-115.
- [18] Awate SA, Patil RB, Ghode PD, Patole V, Pachauri D, Sherief SH. Aphrodisiac activity of aqueous extract of *Glycyrrhiza glabra* in male wistar rats. *World J Pharm Res.* 2012; 1(2): 371-378.
- [19] Arcasoy HB, Erenmemisoglu A, Tekol Y, Kurucu S, Kartal M. Effect of *Tribulus terrestris* L. saponin mixture on some smooth muscle preparations: a preliminary study. *Boll Chim Farm.* 1998; 137(11) 473-475.
- [20] Singh S, Gupta YK. Aphrodisiac activity of *Tribulus terrestris* Linn. in experimental models in rats. *J Men Health's.* 2011; 8(1): S75- S77.
- [21] Hosseinzadeh H, Ziaee T, Sadeghi A. The effect of saffron, *Crocus sativus* stigma, extract and its constituents, safranal and crocin on sexual behaviors in normal male rats. *Phytomedicine.* 2008; 15(6): 491-495.
- [22] Shamsa A, Hosseinzadeh H, Molaei M, Shakeri M T, Rajabi O. Evaluation of *Crocus sativus* L.(saffron) on male erectile dysfunction: a pilot study. *Phytomedicine.* 2008; 16(8):690-693.

- [23] Dhawan K, Kumar S, Sharma A. Aphrodisiac activity of methanol extract of leaves of *Passiflora incarnata* Linn. in mice. *Phytother Res.* 2003; 17(4): 401-403.
- [24] Muanya CA, Odukoya OA. Lipid peroxidation as index of activity in aphrodisiac herbs. *J Plant Sci.* 2008; 3(1): 92-98.
- [25] Baylav N. *Eczacılık Tarihi. Yörük Matbaası, İstanbul.* 1968.
- [26] Nicolas M. Un électuaire de jeunesse et une fête de printemps à Manisa (Turquie). In: *Le Cuisinier et le Philosophe, Hommage à Maxime Rodinson, Maisonneuve et Larose. Paris.* 1982, pp. 71-80.
- [27] Şimsek F. Mithridaticum and Mesir: The Story of an Antidote from Antiquity into Ottoman Times. *MJH.* 2012; 2(2): 243-250.
- [28] Hinneburg I, Dorman HJD, Hiltunen R. Antioxidant activities of extracts from selected culinary herbs and spices. *Food Chem.* 2006; 97(1): 122-129.
- [29] Li HB, Wong CC, Cheng KW, Chen F. Antioxidant properties *in vitro* and total phenolic contents in methanol extracts from medicinal plants. *LWTI.* 2008; 41(3): 385-390.
- [30] Aqil F, Ahmad I, Mehmood Z. Antioxidant and free radical scavenging properties of twelve traditionally used Indian medicinal plants. *Turk J Biol.* 2006; 30: 177-183.
- [31] Wojdyło A, Oszmiański J, Czemerys R. Antioxidant activity and phenolic compounds in 32 selected herbs. *Food Chem.* 2007; 105(3): 940-949.
- [32] Assimopoulou A, Zlatanov S, Papageorgiou V. Antioxidant activity of natural resins and bioactive triterpenes in oil substrates. *Food Chem.* 2005; 92(4): 721-727.
- [33] Güven AN. Master Thesis. Mesir macununun antioksidan aktivitesinin ve reolojik özelliklerinin belirlenmesi. Department of Food Engineering, Faculty of Science, İstanbul Technical University, Maslak, İstanbul, Turkey, 2010.
- [34] Martins N, Barros L, Santos-Buelga C, Ferreira ICFR. Antioxidant potential of two Apiaceae plant extracts: A comparative study focused on the phenolic composition. *Ind Crops Prod.* 2016; 79: 188-194.
- [35] Gülçin İ, Oktay M, Kırıcı E, Küfrevioğlu Öİ. Screening of antioxidant and antimicrobial activities of anise (*Pimpinella anisum* L.) seed extracts. *Food Chem.* 2003; 83(3): 371-382.
- [36] Peschel W, Sánchez-Rabeneda F, Diekmann W, Plescher A, Gartzía I, Jiménez D, Lamuela-Raventós R, Buxaderas S, Codina C. An industrial approach in the search of natural antioxidants from vegetable and fruit wastes. *Food Chem.* 2006; 97(1): 137-150.
- [37] Lee SE, Ju EM, Kim JH. Free radical scavenging and antioxidant enzyme fortifying activities of extracts from *Smilax china* root. *Exp Mol Med.* 2001; 33(4): 263-268.
- [38] Kim KK, Kang YH, Kim DJ, Kim TW, Choe M. Comparison of antioxidant, α -glucosidase inhibition and anti-inflammatory activities of the leaf and root extracts of *Smilax china* L. *J Nutr Health.* 2013; 46 (4): 315-323.
- [39] Burits M, Bucar F. Antioxidant activity of *Nigella sativa* essential oil. *Phytother Res.* 2000; 14(5): 323-328.
- [40] Erkan N, Ayranci G, Ayranci E. Antioxidant activities of rosemary (*Rosmarinus officinalis* L.) extract, blackseed (*Nigella sativa* L.) essential oil, carnosic acid, rosmarinic acid and sesamol. *Food Chem.* 2008; 110(1): 76-82.
- [41] Khalaf NA, Shakya AK, Al-Othman A, El-Agbar Z, Farah H. Antioxidant activity of some common plants. *Turk J Biol.* 2008; 32: 51-55.
- [42] Reddy N, Nagoor Vali D, Rani M, Rani S. Evaluation of antioxidant, antibacterial and cytotoxic effects of green synthesized silver nanoparticles by *Piper longum* fruit. *Mater Sci Eng C Mater Biol App.* 2014; 34: 115-122.
- [43] Škerget M, Kotnik P, Hadolin M, Hraš AR, Simonič M, Kneza Ž. Phenols, proanthocyanidins, flavones and flavonols in some plant materials and their antioxidant activities. *Food Chem.* 2005; 89(2): 191-198.
- [44] Ferreira A, Proença C, Serralheiro MLM, Araújo MEM. The *in vitro* screening for acetylcholinesterase inhibition and antioxidant activity of medicinal plants from Portugal. *J Ethnopharmacol.* 2006; 108(1): 31-37.
- [45] Juntachote T, Berghofer E. Antioxidative properties and stability of ethanolic extracts of Holy basil and Galangal. *Food Chem.* 2005; 92(2): 193-202.
- [46] Mayachiew P, Devahastin S. Antimicrobial and antioxidant activities of Indian gooseberry and galangal extracts. *LWT.* 2008; 41(7): 1153-1159.

- [47] Shen X, Chen W, Zheng Y, Lei X, Tang M, Wang H, Song F. Chemical composition, antibacterial and antioxidant activities of hydrosols from different parts of *Areca catechu* L. and *Cocos nucifera* L. *Ind Crops Prod.* 2017; 96: 110-119.
- [48] Chakraborty M, Mitra A. The antioxidant and antimicrobial properties of the methanolic extract from *Cocos nucifera* mesocarp. *Food Chem.* 2008; 107(3): 994-999.
- [49] Tajuddin, Ahmad S, Latif A, Qasmi IA. Aphrodisiac activity of 50% ethanolic extracts of *Myristica fragrans* Houtt. (nutmeg) and *Syzygium aromaticum* (L) Merr. & Perry (clove) in male mice: a comparative study. *ISCMR.* 2003; 3(6): 1-5.
- [50] Chatterjee S, Niaz Z, Gautam S, Adhikari S, Variyar P, Sharma A. Antioxidant activity of some phenolic constituents from green pepper (*Piper nigrum* L.) and fresh nutmeg mace (*Myristica fragrans*). *Food Chem.* 2007; 101(2): 515-523.
- [51] Singh G, Marimuthu P, de Heluani CS, Catalan C. Antimicrobial and antioxidant potentials of essential oil and acetone extract of *Myristica fragrans* Houtt. (Aril Part). *J Food Sci.* 2005; 70(2): M141-M148.
- [52] Siddhuraju P, Mohan P, Becker K. Studies on the antioxidant activity of Indian Laburnum (*Cassia fistula* L.): a preliminary assessment of crude extracts from stem bark, leaves, flowers and fruit pulp. *Food Chem.* 2002; 79(1): 61-67.
- [53] Luximon-Ramma A, Bahorun T, Soobrattee MA, Aruoma OI. Antioxidant activities of phenolic, proanthocyanidin, and flavonoid components in extracts of *Cassia fistula*. *J Agric Food Chem.* 2002; 50(18): 5042-5047.
- [54] Verma SK, Jain V, Katewa SS. Blood pressure lowering, fibrinolysis enhancing and antioxidant activities of Cardamom (*Elettaria cardamomum*). *Indian J Biochem Biophys.* 2009; 46: 503-506.
- [55] Singh G, Kiran S, Marimuthu P, Isidorov V, Vinogorova V. Antioxidant and antimicrobial activities of essential oil and various oleoresins of *Elettaria cardamomum* (seeds and pods). *J Sci Food Agr.* 2008; 88(2): 280-289.
- [56] Hazra B, Sarkar R, Biswas S, Mandal N. Comparative study of the antioxidant and reactive oxygen species scavenging properties in the extracts of the fruits of *Terminalia chebula*, *Terminalia belerica* and *Embllica officinalis*. *BMC Complement Altern Med.* 2010; 10(20): 1-15.
- [57] Cheng HY, Lin TC, Yu KH, Yang CM, Lin CC. Antioxidant and Free Radical Scavenging Activities of *Terminalia chebula*. *Biol Pharm Bul.* 2003; 26: 1331-1335.
- [58] Vijayakumar RS, Surya D, Nalini N. Antioxidant efficacy of black pepper (*Piper nigrum* L.) and piperine in rats with high fat diet induced oxidative stress. *Redox Rep.* 2004; 9(2): 105-110.
- [59] Gülçin İ. The antioxidant and radical scavenging activities of black pepper (*Piper nigrum*) seeds. *Int J Food Sci Nutr.* 2005; 56(79): 491-499.
- [60] Lee K, Shibamoto T. Antioxidant property of aroma extract isolated from clove buds [*Syzygium aromaticum* (L.) Merr. et Perry]. *Food Chem.* 2001; 74(4): 443-448.
- [61] Atawodi SE, Atawodi, JC, Pfundstein B, Spiegelhalder, Bartsch Hand, Owen R. Assessment of the polyphenol components and *in vitro* antioxidant properties of *Syzygium aromaticum* (L.) Merr.& Perry. *EJEAFChe.* 2011; 10(3): 1970-1978.
- [62] El-Ghorab AH, Nauman M, Anjum FM, Shahzad Hussain S, Nadeem M. A Comparative study on chemical composition and antioxidant activity of ginger (*Zingiber officinale*) and cumin (*Cuminum cyminum*). *J Agric Food Chem.* 2010; 58(14): 8231-8237.
- [63] Rebey IB, Jabri-Karoui I, Hamrouni-Sellami I, Bourgou S, Limam F, Marzouk B. Effect of drought on the biochemical composition and antioxidant activities of cumin (*Cuminum cyminum* L.) seeds. *Ind Crops Prod.* 2012; 36(1): 238-245.
- [64] Ramadan MF, Kroh LW, Mörsel JT. Radical scavenging activity of black cumin (*Nigella sativa* L.), coriander (*Coriandrum sativum* L.), and niger (*Guizotia abyssinica* Cass.) crude seed oils and oil fractions. *J Agric Food Chem.* 2003; 51(24): 6961-6969.
- [65] Samojlik I, Lakić N, Mimica-Dukić N, Đaković-Švajcer K, Božin B. Antioxidant and hepatoprotective potential of essential oils of coriander (*Coriandrum sativum* L.) and caraway (*Carum carvi* L.) (Apiaceae). *J Agric Food Chem.* 2010; 58(15): 8848-8853.
- [66] Roby M, Sarhan M, Selim K, Khalel K. Evaluation of antioxidant activity, total phenols and phenolic compounds in thyme (*Thymus vulgaris* L.), sage (*Salvia officinalis* L.), and marjoram (*Origanum majorana* L.) extracts. *Ind Crops Prod.* 2013; 43: 827-831.

- [67] Economou KD, Oreopoulou V, Thomopoulos CD. Antioxidant activity of some plant extracts of the family Labiatae. *J Am Oil Chem Soc.* 1991; 68 (2): 109-113.
- [68] Visavadiya NP, Soni B, Dalwadi N. Evaluation of antioxidant and anti-atherogenic properties of *Glycyrrhiza glabra* root using *in vitro* models. *Int J Food Sci Nutr.* 2009; 60(2): 135-149.
- [69] Mohamed A, Ali S, EL-Baz F, Hegazy A, Kord M. Chemical composition of essential oil and *in vitro* antioxidant and antimicrobial activities of crude extracts of *Commiphora myrrha* resin. *Ind Crops Prod.* 2014; 57: 10-16.
- [70] Yizhong, Sun M, Xing J, Corke H. Antioxidant phenolic constituents in roots of *Rheum officinale* and *Rubia cordifolia*: Structure–radical scavenging activity relationships. *J Agric Food Chem.* 2004; 52(26): 7884-7890.
- [71] Oktay M, Gülçin İ, Küfrevioğlu Öİ. Determination of *in vitro* antioxidant activity of fennel (*Foeniculum vulgare*) seed extracts. *Food Sci Technol.* 2003; 36(2): 263-271.
- [72] Singh G, Maurya S, de Lampasona M, Catalan C. Chemical constituents, antifungal and antioxidative potential of *Foeniculum vulgare* volatile oil and its acetone extract. *Food Control.* 2006; 17(9): 745-752.
- [73] Karimi E, Oskoueian E, Hendra R, Jaafar HZE. Evaluation of *Crocus sativus* L. Stigma phenolic and flavonoid compounds and its antioxidant activity. *Molecules.* 2010; 15(9): 6244-6256.
- [74] Assimopoulou AN, Sinakos Z, Papageorgiou VP. Radical scavenging activity of *Crocus sativus* L. extract and its bioactive constituents. *Phytother Res.* 2005; 19(11): 997-1000.
- [75] Gardeli C, Vassiliki P, Athanasios M, Kibouris T, Komaitis M. Essential oil composition of *Pistacia lentiscus* L. and *Myrtus communis* L.: Evaluation of antioxidant capacity of methanolic extracts. *Food Chem.* 2008; 107(3): 1120-1130.
- [76] Barra A, Coroneo V, Dessi S, Cabras P, Angioni A. Characterization of the volatile constituents in the essential oil of *Pistacia lentiscus* L. from different origins and its antifungal and antioxidant activity. *J Agric Food Chem.* 2007; 55(17): 7093-7098.
- [77] Palasuwan A, Soogarun S, Lertlum T, Pradnawat P, Wiwanitkit V. Inhibition of Heinz body induction in an *in vitro* model and total antioxidant activity of medicinal Thai plants. *Asian Pacific J Cancer Prev.* 2005; 6: 458-463.
- [78] Laghari AQ, Memon S, Nelofar A, Laghari AH. Extraction, identification and antioxidative properties of the flavonoid-rich fractions from leaves and flowers of *Cassia angustifolia*. *Am J Analyt Chem.* 2011; 2: 871-878.
- [79] Kim JW, Minamikawa T. Hydroxyl radical-scavenging effects of spices and scavengers from brown mustard (*Brassica nigra*). *Biosci Biotechnol Biochem.* 1997; 61(1): 118-123.
- [80] Lin CC, Wu SJ, Chang CH, Ng LT. Antioxidant activity of *Cinnamomum cassia*. *Phytother Res.* 2003; 17(7): 726-730.
- [81] Yang CH, Li RX, Chuang Ly. Antioxidant activity of various parts of *Cinnamomum cassia* extracted with different extraction methods. *Molecules.* 2012; 17(6): 7294-7304.
- [82] Jayaprakasha GK, Rao LJM, Sakariah KK. Volatile constituents from *Cinnamomum zeylanicum* fruit stalks and their antioxidant activities. *J Agric Food Chem.* 2003; 51 (15): 4344-4348.
- [83] Faria A, Oliveira J, Neves P, Gameiro P, Santos-Buelga C, de Freitas V, Mateus N. Antioxidant properties of prepared blueberry (*Vaccinium myrtillus*) extracts. *J Agric Food Chem.* 2005; 53(17): 6896-6902.
- [84] Shyamala BN, Naidu MM, Sulochanamma G, Srinivas P. Studies on the antioxidant activities of natural vanilla extract and its constituent compounds through *in vitro* models. *J Agric Food Chem.* 2007; 55(19): 7738-7743.
- [85] Kamat JP, Ghosh A, Devasagayam T. Vanillin as an antioxidant in rat liver mitochondria: Inhibition of protein oxidation and lipid peroxidation induced by photosensitization. *Mol Cell Biochem.* 2000; 209 (1): 47-53.
- [86] Stoilova I, Krastanov A, Stoyanova A, Denev P, Gargova S. Antioxidant activity of a ginger extract (*Zingiber officinale*). *Food Chem.* 2007; 102(3): 764-770.
- [87] Singh G, Kapoor I, Singh P, Heluani CS, Lampasona MP, Catalan CAN. Chemistry, antioxidant and antimicrobial investigations on essential oil and oleoresins of *Zingiber officinale*. *Food Chem Toxicol.* 2008; 46 (10): 3295-3302.
- [88] Toda S, Miyase T, Arichi H, Tanizawa H, Takino Y. Natural antioxidants.III. Antioxidative components isolated from rhizome of *Curcuma longa* L. *Chem Pharm Bull.* 1985; 33; 1725-1728.
- [89] Ramsewak R, DeWitt D, Nair M. Cytotoxicity, antioxidant and anti-inflammatory activities of Curcumins I–III from *Curcuma longa*. *Phytomedicine.* 2000; / (4): 303-308.
- [90] Mau JL, Lai EYC, Wang NP, Chen CC, Chang CH, Chyau CC. Composition and antioxidant activity of the essential oil from *Curcuma zedoaria*. *Food Chem.* 2003; 82(4): 583-591.

- [91] Özdemir E, Alpınar K. An ethnobotanical survey of medicinal plants in western part of central Taurus Mountains: Aladaglar (Nigde – Turkey). *J Ethnopharmacol.* 2015; 166: 53-65.
- [92] Cakilcioglu U, Khatun S, Turkoglu I, Hayta S. Ethnopharmacological survey of medicinal plants in Maden (Elazığ-Turkey). *J Ethnopharmacol.* 2011; 137: 469-486.
- [93] Ozturk M, Uysal I, Gucl S, Altundag E, Dogan, Baslar S. Medicinal uses of natural dye-yielding plants in Turkey. *RJTA.* 2013; 17(2): 69-80.
- [94] Sezik E, Yeşilada E, Honda G, Takaishi Y, Takeda Y, Tanaka T. Traditional medicine in Turkey IX: Folk medicine in north-west Anatolia. *J Ethnopharmacol.* 1999; 3: 195-210.
- [95] Akbulut S, Bayramoglu MM. The trade and use of some medical and aromatic herbs in Turkey. *Ethno Med.* 2013; 7(2): 67-77.
- [96] Altundag E, Ozturk M. Ethnomedicinal studies on the plant resources of east Anatolia, Turkey. *Procedia Soc Behav Sci.* 2011; 19 :756-777.
- [97] Gürdal B, Kültür Ş. An ethnobotanical study of medicinal plants in Marmaris (Muğla, Turkey). *J Ethnopharmacol.* 2013; 1: 113-126.
- [98] Sargin SA. Ethnobotanical survey of medicinal plants in Bozyazı district of Mersin, Turkey. *J Ethnopharmacol.* 2015; 173: 105-126.
- [99] Sezik E, Yeşilada E, Honda G, Takaishi Y, Takeda Y, Tanaka T. Traditional medicine in Turkey X. Folk medicine in Central Anatolia. *J Ethnopharmacol.* 2001; 2-3: 95-115.
- [100] Tetik F, Civelek S, Cakilcioglu U. Traditional uses of some medicinal plants in Malatya (Turkey). *J Ethnopharmacol.* 2013; 1: 331-346.
- [101] Kaval I, Behçet L, Cakilcioglu U. Ethnobotanical study on medicinal plants in Geçitli and its surrounding (Hakkari-Turkey). *J Ethnopharmacol.* 2014; 1: 171-184.
- [102] Mükemre M, Behçet L, Cakilcioglu U. Ethnobotanical study on medicinal plants in villages of Çatak (Van-Turkey). *J Ethnopharmacol.* 2015; 361-374.
- [103] Polat R, Cakilcioglu U, Satıl F. Traditional uses of medicinal plants in Solhan (Bingöl – Turkey). *J Ethnopharmacol.* 2013; 148: 951-963.
- [104] Güzel Y, Güzelşemme M, Miski M. Ethnobotany of medicinal plants used in Antakya: A multicultural district in Hatay Province of Turkey. *J Ethnopharmacol.* 2015; 174: 118 - 152.
- [105] Hayta S, Polat R, Selvi S. Traditional uses of medicinal plants in Elazığ (Turkey). *J Ethnopharmacol.* 2014; 3: 613-623.
- [106] Polat R, Cakilcioglu U, Kaltalioglu K, Ulsan MD, Türkmen Z. An ethnobotanical study on medicinal plants in Espiye and its surrounding (Giresun-Turkey). *J Ethnopharmacol.* 2015; 163: 1-11.
- [107] Sargin SA, Selvi S, López V. Ethnomedicinal plants of Sarigöl district (Manisa). Turkey. *J Ethnopharmacol.* 2015; 171: 64-84.
- [108] Fakir H, Korkmaz M, Güller B. Medicinal Plant Diversity of Western Mediterranean Region in Turkey. *JABS.* 2009; 3(2): 30-40.
- [109] Kilic O, Bağci E. An ethnobotanical survey of some medicinal plants in Keban (Elazığ-Turkey). *J Med Plants Res.* 2013; 7(23): 1675-1684.
- [110] Akgül G, Yılmaz N, Celep A, Celep F, Çakılıoğlu U. Ethnobotanical purposes of plants sold by herbalists and folk bazaars in the center of Cappadocia (Nevşehir, Turkey). *Indian J Tradit Knowle.* 2016; 15(1): 103-108.

This is an open access article which is publicly available on our journal's website under Institutional Repository at <http://dSPACE.marmara.edu.tr>.